

113068

4 MONTH SUPPLY

PARADEFENSE FOR CATS

For use ONLY on Cats 8 weeks and Older and Weighing 5 to 9 lbs

4 MONTH SUPPLY

4 MONTH SUPPLY

PARADEFENSE FOR CATS

4 MONTH SUPPLY



CONTAINS IMIDACLOPRID AND PYRIPROXYFEN

For use ONLY on Cats 8 weeks and Older and Weighing 5 to 9 lbs

PARADEFENSE FOR CATS

PARADEFENSE FOR CATS

For use ONLY on Cats 8 weeks and Older and Weighing 5 to 9 lbs

For use ONLY on Cats 8 weeks and Older and Weighing 5 to 9 lbs

● KILLS FLEAS AND FLEA EGGS ● PREVENTS REINFESTATION ● FRAGRANCE-FREE



ACTIVE INGREDIENTS:	
Imidacloprid	9.10%
Pyriproxyfen	0.46%
OTHER INGREDIENTS	90.44%
TOTAL	100.00%

KEEP OUT OF REACH OF CHILDREN CAUTION

See back panel for precautionary statements and directions for use.

Convenient, easy-to-apply monthly topical solution. NET CONTENTS: 0.056 fl. oz (1.6ml), Contains 4 tubes, each 0.014 fl.oz (0.4ml)

MADE IN THE U.S.A.



Convenient to use. paraDEFENSE™ for Cats remains effective even after bathing or water immersion. Monthly use of paraDEFENSE™ on cats kills fleas that may cause Flea Allergy Dermatitis.

FIRST AID
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. For medical emergencies or consumer questions call 1-855-844-4375.

IF SWALLOWED: Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.

IF IN EYES: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

IF ON SKIN OR CLOTHING: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment or advice.

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS CAUTION: Harmful if swallowed. Causes moderate eye irritation. Avoid contact with eyes or clothing. Wash hands thoroughly with soap and warm water after handling. Keep out of reach of children. Do not contaminate feed or food.

HAZARDS TO DOMESTIC ANIMALS For external use only. Do not apply to cats or kittens under 8 weeks of age or weighing less than 5 lbs. As with any product, consult your veterinarian before using this product on debilitated, aged, pregnant or nursing cats. Individual sensitivities, while rare, may occur after using ANY pesticide product for cats. If signs persist, or become more severe consult a veterinarian immediately. If your cat is on medication, consult your veterinarian before using this or any other product.

DIRECTIONS FOR USE
It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Do not allow children to apply product.

READ ENTIRE LABEL BEFORE EACH USE

RESTRICTIONS USE ONLY ON CATS OR KITTENS 8 WEEKS AND OLDER WEIGHING 5 to 9 lbs. DO NOT USE ON OTHER ANIMALS

RESTRICTIONS (Cont.)
DO NOT EXCEED LABELED DOSAGE AMOUNT FOR CATS. DO NOT APPLY MORE THAN ONE (1) TUBE PER TREATMENT ON ANY CAT. DO NOT HAVE CONTACT OR ALLOW CHILDREN TO HAVE CONTACT WITH TREATED AREA UNTIL COMPLETELY DRY.

SIDE EFFECTS (CATS): Monitor your cat after application. Side effects, although very rare, may include signs of skin irritation such as redness, scratching, or other signs of discomfort. Gastrointestinal signs such as hypersalivation, vomiting or diarrhea have also been reported. If these or other side effects (such as lethargy) occur, consult your veterinarian or call, 1-855-844-4375.


HOW TO APPLY
1. Remove one applicator tube from the package.
2. Hold applicator tube in an upright position facing away from you and your pet's face and eyes. Pull cap off tube.
3. Turn the cap around and place other end of cap back on tube.
4. Twist cap to break seal, then remove cap from tube.
5. Part the hair on the neck at the base of the skull until the skin is visible. Place the tip of the tube on the skin and squeeze the tube to expel the entire contents directly on the skin. Do not get this product in your pet's eyes, or allow your pet to ingest this product. The product is bitter tasting and salivation may occur for a short time if the pet licks the product immediately after treatment. Treatment at the base of the skull will minimize the opportunity for the cat to lick the product. Do not allow the product to run off.
6. Discard empty tube as described in the Storage and Disposal section.
7. Under normal conditions this product is effective for a month. However, in case of severe flea infestation, retreatment may be necessary earlier than four (4) weeks. On cats, do not retreat more often than once every seven (7) days. After flea control is attained, return to a monthly treatment schedule.

STORAGE AND DISPOSAL

PESTICIDE STORAGE: Store in a cool, dry place inaccessible to children and pets.

PESTICIDE DISPOSAL AND CONTAINER HANDLING: Non-refillable container. If empty: Do not reuse or refill this container. Place in trash or offer for recycling if available. If partly filled: Call your local solid waste agency or 1-888-261-4657 for disposal instructions. Never place unused product down any indoor or outdoor drain.

For more information call our experts at 1-888-261-4657
Manufactured for: CAP IM Supply, Inc. EPA Reg. No. 2596-181-91384
303 Perimeter Center North, Suite 300 EPA Est. No. 2596-OH-1
Atlanta, GA 30346
Made in the USA



RM113068

PARADEFENSE FOR CATS

PARADEFENSE™
FOR CATS



For use
ONLY on Cats
8 weeks and Older
and Weighing
5 to 9 lbs

PARADEFENSE™
FOR CATS

4 MONTH
SUPPLY

PARADEFENSE™
FOR CATS



- KILLS FLEAS AND FLEA EGGS
- PREVENTS REINFESTATION
- FRAGRANCE-FREE




For use
ONLY on Cats
8 weeks and Older
and Weighing
5 to 9 lbs

PARADEFENSE™
FOR CATS

4 MONTH
SUPPLY



RM113504

 <p>The Standard Group 2415 Plantside Drive Louisville, Kentucky 40269-0037 Phone: 502-491-2760 Fax: 502-491-2767 www.thestandardgroup.com</p>	<p>QUALITY ASSURANCE: Our TSG Quality Assurance team has thoroughly reviewed this proof. We present this proof to you for your careful review. Please check to satisfy your requests for all components including the following: composition, color, size, spelling, punctuation, type flow, fonts, UPC, and trapping. All PANTONE colors will be targeted to the PMS BOOK or ink draw downs. Omits to spot coating must be indicated by the signee on the proof before sending the proof back to plant for production.</p>		<p>SFC Approvals</p>					
	Customer <u>Hartz</u>	CLIENT APPROVAL						
Job Description <u>Cat 5-9 lbs.</u>	Signature: _____							
Cad Design# <u>N15-G-09R2</u> CSR <u>Sharon R.</u>	Date: _____							
1st PMS #	2nd BLACK	3rd CYAN	4th MAGENTA	5th PMS #	6th PMS #	7th YELLOW	<p><i>Signed approval releases Graphics into production</i></p>	

FIRST AID

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. For medical emergencies or consumer questions call 1-855-844-4375.

IF SWALLOWED: Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.

IF IN EYES: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

IF ON SKIN OR CLOTHING: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment or advice.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS

CAUTION: Harmful if swallowed. Causes moderate eye irritation. Avoid contact with eyes or clothing. Wash hands thoroughly with soap and warm water after handling. Keep out of reach of children. Do not contaminate feed or food.

HAZARDS TO DOMESTIC ANIMALS

For external use only. Do not apply to cats or kittens under 8 weeks of age or weighing less than 5 lbs. As with any product, consult your veterinarian before using this product on debilitated, aged, pregnant or nursing cats. Individual sensitivities, while rare, may occur after using ANY pesticide product for cats. If signs persist, or become more severe consult a veterinarian immediately. If your cat is on medication, consult your veterinarian before using this or any other product.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Do not allow children to apply product.

READ ENTIRE LABEL BEFORE EACH USE

RESTRICTIONS

USE ONLY ON CATS OR KITTENS 8 WEEKS AND OLDER WEIGHING 5 TO 9 lbs.

DO NOT USE ON OTHER ANIMALS
DO NOT EXCEED LABELED DOSE
AMOUNT FOR CATS

DO NOT APPLY MORE THAN ONE (1)
TUBE PER TREATMENT ON ANY CAT
DO NOT HAVE CONTACT OR ALLOW
CHILDREN TO HAVE CONTACT WITH
TREATED AREA UNTIL COMPLETELY DRY.

RM113498



113498

SIDE EFFECTS (CATS): Monitor your cat after application. Side effects, although very rare, may include signs of skin irritation such as redness, scratching, or other signs of discomfort. Gastrointestinal signs such as hypersalivation, vomiting or diarrhea have also been reported. If these or other side effects (such as lethargy) occur, consult your veterinarian or call, 1-855-844-4375.

HOW TO APPLY

1. Remove one applicator tube from the package.
2. Hold applicator tube in an upright position **facing away from you and your pet's face and eyes**. Pull cap off tube.
3. Turn the cap around and place other end of cap back on tube.
4. Twist cap to break seal, then remove cap from tube.
5. Part the hair on the neck at the base of the skull until the skin is visible. Place the tip of the tube on the skin and squeeze the tube to expel the entire contents directly on the skin. **Do not get this product in your pet's eyes, or allow your pet to ingest this product.** The product is bitter tasting and salivation may occur for a short time if the pet licks the product immediately after treatment. Treatment at the base of the skull will minimize the opportunity for the cat to lick the product. Do not allow the product to run off.
6. Discard empty tube as described in the Storage and Disposal section.
7. Under normal conditions this product is effective for a month. However, in case of severe flea infestation, retreatment may be necessary earlier than four (4) weeks. On cats, do not retreat more often than once every seven (7) days. After flea control is attained, return to a monthly treatment schedule.

STORAGE AND DISPOSAL

PESTICIDE STORAGE: Store in a cool, dry place inaccessible to children and pets.

PESTICIDE DISPOSAL AND CONTAINER HANDLING:


Non-refillable container. **If empty:** Do not reuse or refill this container. Place in trash or offer for recycling if available. **If partly filled:** Call your local solid waste agency or 1-888-261-4657 for disposal instructions. Never place unused product down any indoor or outdoor drain.

For more information call our experts at 1-888-261-4657

Manufactured for: CAP IM Supply, Inc.
303 Perimeter Center North, Suite 300
Atlanta, GA 30346
Made in the USA

EPA Reg. No. 2596-181-91384
EPA Est. No. 2596-OH-1

Assumes Full Coverage at the center of the tube
May overlap at the open end.



PARADEFENSE™

FOR CATS

CONTAINS IMIDACLOPRID AND PYRIPROXYFEN

REMINDER STICKERS

THE FIRST TIME YOU TREAT YOUR CAT, PLACE THE FIRST MONTH STICKER ON YOUR CALENDAR.
APPLY PARA|DEFENSE™ MONTHLY, EVERY 30 DAYS. USE THESE CALENDAR STICKERS
TO REMIND YOU WHEN YOUR CAT IS DUE FOR THE NEXT APPLICATION.



For more information call our experts at 1-888-261-4657
Manufactured for: CAP IM Supply, Inc.
303 Perimeter Center North, Suite 300 Atlanta, GA 30346

403077
MADE IN THE U.S.A.

TRIM: 2.5" X 3.25"
COLOR: CMYK424 DATE: 12/20/15
Hartz_Cats_Stickers


digitalcolor
job 152471



RM113069

4 MONTH SUPPLY

4 MONTH SUPPLY

4 MONTH SUPPLY

4 MONTH SUPPLY

PARADEFENSE™ FOR CATS

PARADEFENSE™ FOR CATS

PARADEFENSE™ FOR CATS

PARADEFENSE™ FOR CATS

CONTAINS IMIDACLOPRID AND PYRIPROXYFEN



For use ONLY on Cats 8 Weeks and Older and Weighing **over 9 lbs**

For use ONLY on Cats 8 weeks and Older and Weighing **over 9 lbs**

For use ONLY on Cats 8 weeks and Older and Weighing **over 9 lbs**

For use ONLY on Cats 8 weeks and Older and Weighing **over 9 lbs**

- KILLS FLEAS AND FLEA EGGS
- PREVENTS REINFESTATION
- FRAGRANCE-FREE



ACTIVE INGREDIENTS:	
Imidacloprid	9.10%
Pyriproxyfen	0.46%
OTHER INGREDIENTS	90.44%
TOTAL	100.00%

KEEP OUT OF REACH OF CHILDREN CAUTION

See back panel for precautionary statements and directions for use.

Convenient, easy-to-apply monthly topical solution.
NET CONTENTS: 0.108 fl. oz (3.2ml), Contains 4 tubes, each 0.027 fl.oz (0.8ml)

MADE IN THE U.S.A.



Convenient to use. paraDEFENSE™ for Cats remains effective even after bathing or water immersion. Monthly use of paraDEFENSE™ on cats kills fleas that may cause Flea Allergy Dermatitis.

FIRST AID
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. For medical emergencies or consumer questions call 1-855-844-4375.

IF SWALLOWED: Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.

IF IN EYES: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

IF ON SKIN OR CLOTHING: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment or advice.

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS CAUTION: Harmful if swallowed. Causes moderate eye irritation. Avoid contact with eyes or clothing. Wash hands thoroughly with soap and warm water after handling. Keep out of reach of children. Do not contaminate feed or food.

HAZARDS TO DOMESTIC ANIMALS For external use only. Do not apply to cats or kittens under 8 weeks of age or weighing less than 9 lbs. As with any product, consult your veterinarian before using this product on debilitated, aged, pregnant or nursing cats. Individual sensitivities, while rare, may occur after using ANY pesticide product for cats. If signs persist, or become more severe consult a veterinarian immediately. If your cat is on medication, consult your veterinarian before using this or any other product.

DIRECTIONS FOR USE
It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Do not allow children to apply product.

READ ENTIRE LABEL BEFORE EACH USE

RESTRICTIONS USE ONLY ON CATS OR KITTENS 8 WEEKS AND OLDER WEIGHING OVER 9 lbs. DO NOT USE ON OTHER ANIMALS

RESTRICTIONS (Cont.)
DO NOT EXCEED LABELED DOSAGE AMOUNT FOR CATS. DO NOT APPLY MORE THAN ONE (1) TUBE PER TREATMENT ON ANY CAT. DO NOT HAVE CONTACT OR ALLOW CHILDREN TO HAVE CONTACT WITH TREATED AREA UNTIL COMPLETELY DRY.

SIDE EFFECTS (CATS): Monitor your cat after application. Side effects, although very rare, may include signs of skin irritation such as redness, scratching, or other signs of discomfort. Gastrointestinal signs such as hypersalivation, vomiting or diarrhea have also been reported. If these or other side effects (such as lethargy) occur, consult your veterinarian or call, 1-855-844-4375.

- HOW TO APPLY**
1. Remove one applicator tube from the package.
 2. Hold applicator tube in an upright position **facing away from you and your pet's face and eyes.** Pull cap off tube.
 3. Turn the cap around and place other end of cap back on tube.
 4. Twist cap to break seal, then remove cap from tube.
 5. Part the hair on the neck at the base of the skull until the skin is visible. Place the tip of the tube on the skin and squeeze the tube to expel the entire contents directly on the skin. *Do not get this product in your pet's eyes, or allow your pet to ingest this product. The product is bitter tasting and salivation may occur for a short time if the pet licks the product immediately after treatment.* Treatment at the base of the skull will minimize the opportunity for the cat to lick the product. Do not allow the product to run off.
 6. Discard empty tube as described in the Storage and Disposal section.
 7. Under normal conditions this product is effective for a month. However, in case of severe flea infestation, retreatment may be necessary earlier than four (4) weeks. On cats, do not retreat more often than once every seven (7) days. After flea control is attained, return to a monthly treatment schedule.

STORAGE AND DISPOSAL
PESTICIDE STORAGE: Store in a cool, dry place inaccessible to children and pets.

PESTICIDE DISPOSAL AND CONTAINER HANDLING: Non-refillable container. If empty: Do not reuse or refill this container. Place in trash or offer for recycling if available. If partly filled: Call your local solid waste agency or 1-888-261-4657 for disposal instructions. Never place unused product down any indoor or outdoor drain.

For more information call our experts at 1-888-261-4657
Manufactured for: CAP IM Supply, Inc. EPA Reg. No. 2596-181-91384
303 Perimeter Center North, Suite 300 EPA Est. No. 2596-OH-1
Atlanta, GA 30346
Made in the USA



PARADEFENSE™
FOR CATS



For use
ONLY on Cats
8 weeks and Older
and Weighing
over 9 lbs

PARADEFENSE™
FOR CATS

4!!!!
MONTH
SUPPLY

PARADEFENSE™
FOR CATS



● KILLS FLEAS AND FLEA EGGS ● PREVENTS REINFESTATION ● FRAGRANCE-FREE



For use
ONLY on Cats
8 weeks and Older
and Weighing
over 9 lbs

PARADEFENSE™
FOR CATS

4!!!!
MONTH
SUPPLY



RM113505



The Standard Group
2415 Plantside Drive
Louisville, Kentucky 40269-0037
Phone: 502-491-2760 Fax: 502-491-2767
www.thestandardgroup.com

QUALITY ASSURANCE: Our TSG Quality Assurance team has thoroughly reviewed this proof. We present this proof to you for your careful review. Please check to satisfy your requests for all components including the following: composition, color, size, spelling, punctuation, type flow, fonts, UPC, and trapping. All PANTONE colors will be targeted to the PMS BOOK or ink draw downs. Omits to spot coating must be indicated by the signee on the proof before sending the proof back to plant for production.

Customer Hartz
Job Description Large Cat
Cad Design# N15-G-09R2 CSR Sharon R.

CLIENT APPROVAL
Signature: _____
Date: _____

1st PMS # 2nd BLACK 3rd CYAN 4th MAGENTA 5th PMS # 6th PMS # 7th YELLOW

*Signed approval releases
Graphics into production*

SFC
Approvals

FIRST AID

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. For medical emergencies or consumer questions call 1-855-844-4375.

IF SWALLOWED: Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.

IF IN EYES: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

IF ON SKIN OR CLOTHING: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment or advice.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS

CAUTION: Harmful if swallowed. Causes moderate eye irritation. Avoid contact with eyes or clothing. Wash hands thoroughly with soap and warm water after handling. Keep out of reach of children. Do not contaminate feed or food.

HAZARDS TO DOMESTIC ANIMALS

For external use only. Do not apply to cats or kittens under 8 weeks of age or weighing less than **9 lbs.** As with any product, consult your veterinarian before using this product on debilitated, aged, pregnant or nursing cats. Individual sensitivities, while rare, may occur after using ANY pesticide product for cats. If signs persist, or become more severe consult a veterinarian immediately. If your cat is on medication, consult your veterinarian before using this or any other product.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Do not allow children to apply product.

READ ENTIRE LABEL BEFORE EACH USE

RESTRICTIONS

USE ONLY ON CATS OR KITTENS 8 WEEKS AND OLDER WEIGHING OVER 9 lbs.

**DO NOT USE ON OTHER ANIMALS
DO NOT EXCEED LABELED DOSAGE
AMOUNT FOR CATS**

**DO NOT APPLY MORE THAN ONE (1)
TUBE PER TREATMENT ON ANY CAT
DO NOT HAVE CONTACT OR ALLOW
CHILDREN TO HAVE CONTACT WITH
TREATED AREA UNTIL COMPLETELY DRY.**

RM113499



113499

SIDE EFFECTS (CATS): Monitor your cat after application. Side effects, although very rare, may include signs of skin irritation such as redness, scratching, or other signs of discomfort.

Gastrointestinal signs such as hypersalivation, vomiting or diarrhea have also been reported. If these or other side effects (such as lethargy) occur, consult your veterinarian or call, 1-855-844-4375.

HOW TO APPLY

1. Remove one applicator tube from the package.
2. Hold applicator tube in an upright position **facing away from you and your pet's face and eyes.** Pull cap off tube.
3. Turn the cap around and place other end of cap back on tube.
4. Twist cap to break seal, then remove cap from tube.
5. Part the hair on the neck at the base of the skull until the skin is visible. Place the tip of the tube on the skin and squeeze the tube to expel the entire contents directly on the skin. **Do not get this product in your pet's eyes, or allow your pet to ingest this product.** The product is bitter tasting and salivation may occur for a short time if the pet licks the product immediately after treatment. Treatment at the base of the skull will minimize the opportunity for the cat to lick the product. Do not allow the product to run off.
6. Discard empty tube as described in the Storage and Disposal section.
7. Under normal conditions this product is effective for a month. However, in case of severe flea infestation, retreatment may be necessary earlier than four (4) weeks. On cats, do not retreat more often than once every seven (7) days. After flea control is attained, return to a monthly treatment schedule.

STORAGE AND DISPOSAL

PESTICIDE STORAGE: Store in a cool, dry place inaccessible to children and pets.

PESTICIDE DISPOSAL AND CONTAINER HANDLING:


Non-refillable container. **If empty:** Do not reuse or refill this container. Place in trash or offer for recycling if available. **If partly filled:** Call your local solid waste agency or 1-888-261-4657 for disposal instructions. Never place unused product down any indoor or outdoor drain.

For more information call our experts at 1-888-261-4657

Manufactured for: CAP IM Supply, Inc.
303 Perimeter Center North, Suite 300
Atlanta, GA 30346
Made in the USA

EPA Reg. No. 2596-181-91384
EPA Est. No. 2596-OH-1

Assumes Full Coverage at the center of the tube
May overlap at the open end.



PARADEFENSE™ FOR CATS

CONTAINS IMIDACLOPRID AND PYRIPROXYFEN

REMINDER STICKERS

THE FIRST TIME YOU TREAT YOUR CAT, PLACE THE FIRST MONTH STICKER ON YOUR CALENDAR.
APPLY PARA|DEFENSE™ MONTHLY, EVERY 30 DAYS. USE THESE CALENDAR STICKERS
TO REMIND YOU WHEN YOUR CAT IS DUE FOR THE NEXT APPLICATION.



For more information call our experts at 1-888-261-4657
Manufactured for: CAP IM Supply, Inc.
303 Perimeter Center North, Suite 300 Atlanta, GA 30346

403077
MADE IN THE U.S.A.

TRIM: 2.5" X 3.25"
COLOR: CMYK424 DATE: 12/20/15
Hartz_Cats_Stickers


digitalcolor
job 152471