

CAUTION: Harmful if swallowed. Causes moderate eye irritation. Avoid contact with eyes or clothing. Wash hands thoroughly with soap and warm water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet.

HAZARDS TO DOMESTIC ANIMALS

For external use only. Do not use on puppies under 7 weeks of age or weighing less than 3 lbs. As with any product, consult your veterinarian before using this product on debilitated, aged, pregnant or nursing dogs. Individual sensitivities, while rare, may occur after using ANY pesticide product for dogs. If signs persist, or become more severe, consult a veterinarian immediately. If your dog is on medication, consult your veteringrign before using this or any other product.

FIRST AID

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. For medical emergencies or consumer questions call 1-855-844-4375.

SWALLOWED: Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.

IF IN EYES: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eve. Call a poison control center or doctor for treatment advice.

IF ON SKIN OR CLOTHING: Take off contaminated dothing Rinse skin immediately with plenty of water for 15–20 minutes. Call a poison control center or doctor for treatment or advice.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Do not allow children to apply product.

READ ENTIRE LABEL BEFORE EACH USE

RESTRICTIONS

LISE ONLY ON DOGS OR ON PLIPPIES 7 WEEKS OF AGE OR OLDER WEIGHING 3 to 10 lbs. DO NOT USE ON OTHER ANIMALS DO NOT EXCEED LABELED DOSAGE AMOUNT RM103691 FOR DOGS DO NOT APPLY MORE THAN ONE (1) TUBE PFR TRFATMENT DO NOT HAVE CONTACT OR ALLOW CHILDREN TO HAVE CONTACT WITH TREATED AREA UNTIL COMPLETELY DRY

Side Effects: Monitor your dog after application. Side effects, although very rare, may include signs of skin irritation such as redness, scratching, or other signs of discomfort. Gastrointestinal signs such as vomiting or diarrhea have also been reported. If these or other side effects (such as lethargy) occur, consult your veterinarian or call 1-855-844-4375



DO NOT LISE ON CATS.

Keep cats away from treated dogs for 24 hours. If applied to a cat or indested by a cat, contact your veteringrign immediately

HOW TO APPLY

- Remove one applicator tube from the package.
- 2. Hold applicator tube in an upright position facing away from you and your pet's face and eyes. Pull cap off tube.
- furn the cap around and place other end of cap back on tube.
 Twist cap to break seal, then remove cap from tube.
- The dog should be standing for easy application. Part the hair on the dog's back, between the shoulder blades, until the skin is visible. Place the tip of the tube on the skin and squeeze the tube to expel the entire contents directly on the skin. Do not get this product in your dog's eyes or allow your dog to ingest this product. Do not allow the product to run off.
- 6. Discard empty tube as described in the Storage and Disposal section.
- Under normal conditions the product is effective for a month. However, in case of severe flea infestation, retreatment may be necessary earlier than four weeks. Do not re-treat more often than once every seven (7) days. After flea control is obtained, return to a monthly retreatment schedule.

STORAGE AND DISPOSAL

PESTICIDE STORAGE: Store in a cool, dry place inaccessible to children and pets.

PESTICIDE DISPOSAL AND CONTAINER HANDLING:

Non-refillable container. If empty: Do not reuse or refill this container. Place in trash or offer for recycling if available. If partly filled: Call your local solid waste agency or 1-888-845-4921 for disposal instructions. Never place unused product down any indoor or outdoor drain

For more information call our experts at 1-888-845-4921

Manufactured for: CAP IM Supply, Inc. 2825 E. Cottonwood Pkwy. Suite 500 Salt Lake City, UT 84121 Made in the USA

EPA Reg. No. 2596-180-91384 EPA Est. No. 2596-0H-1



EPA Reg. No. 2596-081-384 EPA Reg. No. 2596-081-384 REFORE USE KEAD ENTIRE LABEL NOITUAD OF CHILDREN KEEP OUT OF REACH 0.46% PYRIPROXYFEN **UINYOLDACION WOLG** Only for Use on Dogs and Puppies 7 Weeks or Older Weighing 3 to 10 lbs.-0.014 fl. oz. (0.4 ml) RM113236



Manufactured for: CAP IM Supply, Inc. 2825 E. Cottonwood Pkwy, Suite 500, Salt Lake City, UT 84121 Made in the USA

For more information call our experts at 1-888-845-4921



CAUTION: Harmful if swallowed. Causes moderate ever irritation. Avoid contact with eyes or clothina, Wash hands thoroughly with soap and warm water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet.

HAZARDS TO DOMESTIC ANIMALS

For external use only. Do not use on puppies under 7 weeks of age or weighing less than 11 lbs. As with any product, consult your veterinarian before using this product on debilitated, aged, pregnant or nursing dogs. Individual sensitivities, while rare, may occur after using AIV pestidde product for dogs. It signs persist, or become more severe, consult a veterinarian immediately. If your dog is on medication, consult your veterinarian before using this or any ofther product.

FIRST AID

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. For medical emergencies or consumer questions call 1-855-844-4375.

IF SWALLOWED: Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.

IF IN EYES: Hold eye open and rinse slowly and gently with water for 15–20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

IF ON SKIN OR CLOTHING: Take off contaminated dothing. Rinse skin immediately with plenty of water for 15–20 minutes. Call a poison control center or doctor for treatment or advice.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Do not allow children to apply product.

READ ENTIRE LABEL BEFORE EACH USE

RESTRICTIONS

USE ONLY ON DOGS OR ON PUPPIES 7 WEEKS OF AGE OR OLDER WEIGHING 11 to 20 lbs. Do NOT USE ON OTHER ANIMALS DO NOT TEXPENDATION OF THE ANIMALS DO NOT TAPPLY MORE THAN ONE (1) TUBE PER TREATMENT DO NOT HAVE CONTACT OR ALLOW CHILDREN TO HAVE CONTACT WITH TREATED AREA UNTIL COMPLETELY DRY Side Effects: Monitor your dog after application. Side effects, although very rare, may include signs of skin irritation such as redness, scratching, or other signs of disconfort. Gastromitestical signs such as vomiting or diarrhea have also been reported. If these or other side effects (such as leithargy) occur, consult your veterinarian or call 1-825 844-4375.



DO NOT USE ON CATS.

Keep cats away from treated dogs for 24 hours. If applied to a cat or ingested by a cat, contact your veterinarian immediately.

HOW TO APPLY

- 1. Remove one applicator tube from the package.
- Hold applicator tube in an upright position facing away from you and your pet's face and eyes. Pull cap off tube.
- 3. Turn the cap around and place other end of cap back on tube.
- 4. Twist cap to break seal, then remove cap from tube.
- 5. The day should be standing for easy application. Part the hair on the day shack, between the shoulder blades, until the skin is visible. Place the tip of the tube on the skin and squeeze the tube to expel the entire contents directly on the skin. Do not get this product in your day's eyes or allow your day to ingest this product. Do not allow the product to run off.
- 6. Discard empty tube as described in the Storage and Disposal section.
- Under normal conditions the product is effective for a month. However, in case of severe flea infestation, retreatment may be necessary earlier than four weeks. Do not re-treat more often than once every seven (7) days. After flea control is obtained, return to a monthly retreatment schedule.

STORAGE AND DISPOSAL

PESTICIDE STORAGE: Store in a cool, dry place inaccessible to children and pets.


PESTICIDE DISPOSAL AND CONTAINER HANDLING:

Non-refiltable container, **If empty:** Do not reuse or refilt this container. Place in trash or offer for recycling if available. **If partly filted:** Call your local sold waste agency or 1-888-845-4921 for disposal instructions. Never place unused product down any indoor or outdoor drain.

For more information call our experts at 1-888-845-4921

Manufactured for: CAP IM Supply, Inc. 2825 E. Cottonwood Pkwy, Suite 500 Salt Lake City, UT 84121 Made in the USA

EPA Reg. No. 2596-180-91384 EPA Est. No. 2596-0H-1







Manufactured for: CAP IM Supply, Inc. 2825 E. Cottonwood Pkwy, Suite 500, Salt Lake City, UT 84121 Made in the USA

For more information call our experts at 1-888-845-4921



CAUTION: Harmful if swallowed. Causes moderate eye irritation. Avoid contact with eves or clothing. Wash hands thoroughly with soap and warm water after handling and before eating. drinking, chewing gum, using tobacco or using the toilet

HAZARDS TO DOMESTIC ANIMALS

For external use only. Do not use on puppies under 7 weeks of age or weighing less than 21 lbs. As with any product, consult your veterinarian before using this product on debilitated, aged, pregnant or nursing dogs. Individual sensitivities, while rare, may occur after using ANY pesticide product for dogs. If signs persist, or become more severe, consult a veterinarian immediately. If your dog is on medication, consult your veteringrign before using this or any other product.

FIRST AID

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. For medical emergencies or consumer questions call 1-855-844-4375.

IF SWALLOWED: Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.

IF IN EYES: Hold eve open and rinse slowly and aently with water for 15–20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice

IF ON SKIN OR CLOTHING: Take off contaminated clothing Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment or advice.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Do not allow children to apply product.

READ ENTIRE LABEL BEFORE EACH LISE

RESTRICTIONS

USE ONLY ON DOGS OR ON PUPPIES 7 WEEKS OF AGE OR OLDER WEIGHING 21 to 55 lbs DO NOT LISE ON OTHER ANIMALS RM103693 DO NOT FXCFFD I ABFI FD DOSAGF AMOUNT FOR DOGS DO NOT APPLY MORE THAN ONE (1) TUBE PFR TRFATMENT DO NOT HAVE CONTACT OR ALLOW CHILDREN TO HAVE CONTACT WITH TREATED AREA UNTIL COMPLETELY DRY

Side Effects: Monitor your dog after application. Side effects, although very rare, may include signs of skin irritation such as redness, scratching, or other signs of discomfort. Gastrointesting signs such as vomiting or diarrhea have also been reported. If these or other side effects (such as lethargy) occur, consult your veterinarian or call 1-855-844-4375.



DO NOT USE ON CATS.

Keep cats away from treated doas for 24 hours. If applied to a cat or inaested by a cat, contact vour veterinarian immediately.

HOW TO APPLY

- Remove one applicator tube from the package.
 Hold applicator tube in an upright position facing away from you and your pet's face and eyes, Pull cap off tube.
- 3. Turn the cap around and place other end of cap back on tube.
- 4. Twist cap to break seal, then remove cap from tube.
- 5. The dog should be standing for easy application. Apply the entire contents of the tube evenly to three or four spots on the top of the back from the shoulder to the base of the tail. At each spot, part the hair until the skin is visible. Place the tip of the tube on the skin and gently squeeze to expel a portion of the solution on the skin. Do not apply an excessive amount of solution at any one spot that could cause some of the solution to run off the side of the doa. Do not get this product in your dog's eyes or allow your dog to inaest this product.
- 6. Discard empty tube as described in the Storage and Disposal section.
- 7. Under normal conditions the product is effective for a month. However, in case of severe flea infestation, retreatment may be necessary earlier than four weeks. Do not re-treat more often than once every seven (7) days. After flea control is obtained, return to a monthly retreatment schedule.

STORAGE AND DISPOSAL

PESTICIDE STORAGE: Store in a cool, dry place inaccessible to children and pets.


PESTICIDE DISPOSAL AND CONTAINER HANDLING:

Non-refillable container. If empty: Do not reuse or refill this container. Place in trash or offer for recycling if available. If partly filled: Call your local solid waste agency or 1-888-845-4921 for disposal instructions. Never place unused product down any indoor or outdoor drain.

For more information call our experts at 1-888-845-4921

Manufactured for: CAP IM Supply, Inc. 2825 E. Cottonwood Pkwy, Suite 500 Salt Lake City, UT 84121 Made in the USA

EPA Reg. No. 2596-180-91384 FPA Est. No. 2596-0H-1







Manufactured for: CAP IM Supply, Inc. 2825 E. Cottonwood Pkwy, Suite 500, Salt Lake City, UT 84121 Made in the USA

For more information call our experts at 1-888-845-4921



Side Effects: Monitor your dog after application. Side effects, although very rare, may include signs of skin irritation such as redness, scratching, or other signs of discomfort. Gastrointestinal signs such as vomiting or diarrhea have also been reported. If these or other side effects (such as lethargy) occur, consult your veterinarian or call 1-855-844-4375.



HOW TO APPLY

How to APTLI Remve one applicator tube from the package. Hold applicator tube in an upright position facing away from you and your pet's face and eyes. Pull cap off tube. Jum the cap around not place table rend of cap back on tube. Twist cap to break seal, then remove cap from tube.

- The dog should be standing for easy application. Apply the entire contents of the The eventy to three or four spots on the top of the back from the shoulder to the base of the tail. At each spot, part the hair until the skin is visible. Place the tip of the tube on the skin and gently squeeze to expel a portion of the solution on the skin. Do not apply an excessive amount of solution at any one spot that could cause some of the solution to run off the side of the dog. Do not get this product

in your dag's eyes or allow your dag to ingest this product. Discard empty tube as described in the Storage and Disposal section. Under normal conditions the product is effective for a month. However, in case of severe flea infestation, retreatment may be necessary earlier than four weeks. Do not re-treat more often than once every seven (7) days. After flea control is obtained, return to a monthly retreatment schedule.

STORAGE AND DISPOSAL

PESTICIDE STORAGE: Store in a cool, dry place inaccessible to children and pets.

PESTICIDE DISPOSAL AND CONTAINER HANDLING:

Non-refillable container. If empty: Do not reves or refill this container. Place in trash or offer for recycling if available. If partly filled: Call your local solid waste agency or 1-888-845-4921 for disposal instructions. Never place unused product down any indoor or outdoor drain.

For more information call our experts at 1-888-845-4921



RM113251

CAUTION: Harmful if swallowed. Causes moderate eye irritation. Avoid contact with eyes or clothing. Wash hands thoroughly with soap and warm water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet

HAZARDS TO DOMESTIC ANIMALS

For external use only. Do not use on puppies under 7 weeks of age or weighing less than 55 lbs. As with any product, consult your veterinarian before using this product on debilitated, aged, pregnant or nursing dogs. Individual sensitivities, while rare, may occur after using ANY pesticide product for dogs. If signs persist, or become more severe, consult a veterinarian immediately. If your doa is on medication, consult your veterinarian before using this or any other product.

FIRST AID

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. For medical emergencies or consumer questions call 1-855-844-4375.

IF SWALLOWED: Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.

IF IN EYES: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice

IF ON SKIN OR CLOTHING: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15–20 minutes. Call a poison control center or doctor for treatment or advice.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Do not allow children to apply product.

READ ENTIRE LABEL BEFORE EACH USE

RESTRICTIONS

USE ONLY ON DOGS OR ON PUPPIES 7 WEEKS OF AGE OR OLDER OVER 55 lbs

DO NOT LISE ON OTHER ANIMALS DO NOT EXCEED LABELED DOSAGE AMOUNT

RM113245 FOR DOGS DO NOT APPLY MORE THAN ONE (1) TUBE

PER TREATMENT DO NOT HAVE CONTACT OR ALLOW CHILDREN TO HAVE CONTACT WITH TREATED AREA UNTIL COMPLETELY DRY

to inaest this product. Discard empty tube as described in the Storage and

- Disposal section.
- 7. Under normal conditions the product is effective for a month. However, in case of severe flea infestation, retreatment may be necessary earlier than four weeks. Do not re-treat more often than once every seven (7) days. After flea control is obtained. return to a monthly retreatment schedule.

STORAGE AND DISPOSAL

PESTICIDE STORAGE: Store in a cool, dry place inaccessible to children and pets.

PESTICIDE DISPOSAL AND CONTAINER HANDLING:

Non-refillable container. If empty: Do not reuse or refill this container. Place in trash or offer for recycling if available. If partly filled: Call your local solid waste agency or 1-888-845-4921 for disposal instructions. Never place unused product down any indoor or outdoor drain.

For more information call our experts at 1-888-845-4921

Manufactured for: CAP IM Supply, Inc. 303 Perimeter Center North, Suite 300 Atlanta, GA 30346 Made in the USA

EPA Reg. No. 2596-180-91384 EPA Est. No. 2596-0H-1

your DO NOT USE ON CATS.

1-855-844-4375.

Side Effects: Monitor your dog after application. Side effects. although very rare, may include signs of skin irritation such as

> Keep cats away from treated dogs for 24 hours. If applied to a cat or ingested by a cat, contact vour veterinarian immediately.

redness, scratching, or other signs of discomfort.

Gastrointestinal signs such as vomiting or diarrhea have also been reported. If these or other side effects (such as lethargy) occur,

veteringrign or


(nn)


HOW TO APPLY

Remove one applicator tube from the package.

consult

- 2. Hold applicator tube in an upright position facing away from vou and your pet's face and eyes. Pull cap off tube.
- 3. Turn the cap around and place other end of cap back on tube.
- Twist cap to break seal, then remove cap from tube.
- 5. The dog should be standing for easy application. Apply the entire contents of the tube evenly to three or four spots on the top of the back from the shoulder to the base of the tail. At each spot, part the hair until the skin is visible. Place the tip of the tube on the skin and gently squeeze to expel a portion of the solution on the skin. Do not apply an excessive amount of solution at any one spot that could cause some of the solution to run off the side of the dog. Do not get this product in your dog's eyes or allow your dog







Manufactured for: CAP IM Supply, Inc. 303 Perimeter Center North, Suite 300, Atlanta, GA 30346

Made in the USA

For more information call our experts at 1-888-845-4921