













# SENTRY ADVANCE GUARD2

**FOR DOGS** 

## **SMALL DOG** 3-10 LBS

## **Starts working** through contact

Once-A-Month topical

treatment for fleas on dogs **ONLY for dogs and puppies** 7 weeks or older and weighing 3-10 lbs.

Active Ingredients

Imidacloprid. Pyriproxyfen..... Other ingredients... 0.46% 90.44% .100.00%

KEEP OUT OF REACH OF CHILDREN CAUTION SEE BACK PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS

"SENTRY Advance Guard?" for Dogston on the chartered or distributed by Bayer Hade care LLC Temporary of Advance of

NET CONTENTS: 0.056 fl oz (1.6 mL) 4-9.014 fl oz (0.4 mL) applicators

Contains imidacloprid & pyriproxyfen, the active ingredients used in Advantage® II\*



## STORAGE AND DISPOSAL

PESTICIDE STORAGE: Do not remove tube from the pack until ready to use. Store only in original container in a cool (below 77%-f/25°C), dry place inaccessible to children and pets. Do not refrigerate. Protect from direct sunlight

CONTAINER DISPOSAL: If empty: Non-refillable container. Do not reuse or refill this container. Offer for recycling if available. If partly filled: Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

# SENTRY ADVANCE GUARD2 FOR DOGS

Once per month topical fiea treatment for dogs

This product is for the treatment and prevention of fleas on dogs

Controls multi-stages of flea lifecycle

Kills fleas within 12 hours of andication on doos.

#### READ ENTIRE LARFL REPORT FACH LISE. USE ONLY ON DOGS AND PUPPIES 7 WEEKS OF AGE OR OLDER. PRECAUTIONARY STATEMENTS

HAZAROS TO HUMANS. CAUTION. Harmful if swallowed. Causes moderate eye irritation. Avoid contact with eyes or clothing. Wash hands thoroughly with soap and warm water after handling and before eating, drinking, chewing gurn, using tobacco, or using the toilet.

HAZARDS TO DOMESTIC ANIMALS. For external use on doos only, individual sensitivities. while rare, may occur after using ANY pesticide product for dogs. Do not treat your dog with more than one pesticide product at a time. Over dosing your dog can result in serious illness and even death. Consult a veterinarian before using this product on medicated, debilitated, aged, pregnant or nursing dogs.

FIRST AID If swallowed: • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swalkow. • Do not induce vorniting unless told to do so by the poison control center or doctor. • Do not give anything by mouth to an unconscious person If In eyes: • Hold eye open and rinse slowly and gently with water for 15-20 minutes Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.

If on skin or obtaining: \* Take off contaminated dothing. Rines skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for heatment advice. HOT LINE NUMBER: Have the product container or label with you when calling a poison

control center or doctor or going for treatment. You may also contact 1-800-781-4738 for emergency medical treatment information.

NOTE TO PHYSICIAN: Treat the patient symptomatically.

#### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. TO PREVENT HARM TO YOU AND YOUR DOG, READ ENTIRE LABEL BEFORE EACH USE. FOLLOW ALL DIRECTIONS AND PRECAUTIONARY STATEMENTS CAREFULLY.

- DO NOT ALLOW CHILDREN TO APPLY PRODUCT.
- Do not use on animals other than doos. Do not allow your dog to ingest this product.
- Do not reacoly for 4 weeks.
- rtents of tube on each dog. Do not split one tube between dogs. Do not use multiple tubes on one doa.
- Weigh your dog to be sure you are applying the right dose formulated for the weight of your dog.
   Separate the treated dog from all other dogs and cats for 24 hours after treatment has been applied.
- Do not have contact or allow children to have contact with the treated area until completely dry.

### How to Open:

- 1. Remove product tubes from the package
- 2. Separate one tube from the others. Hold the tube with notched end pointing up and away from the face and body. Use scissors to cut off the narrow end at the notches along the line.
- How to Apply: 1, Invert tube over dog and use open end to part dog's hair.
- 2. Squeeze tube firmly to apply all of the solution to the dog's skin as a spot to the dog's back between the shoulder blades.

Only For Dogs Weighing 3-10 lbs. Do not apply to dogs weighing less than 3 lbs. Apply one tube 0.4 mL (0.014 fi oz) directly to skin as a spot to the dog's back between the shoulde

#### FREQUENCY OF APPLICATION

Use this product monthly for control of flea infestations. Studies show that this product kills fleas within 12 hours of application and lasts for up to 4 weeks. This product kills reintesting fless within 2 hours and prevents further infestations of fless. Apply monthly for fles control. This product is roof and remains effective, even after bothing, swimming, or exposure to surlight or rain.

Side Effects: Monitor your dog after application. Some dogs may experience temporary startle effects when any product is applied Dogs may experience some temporary irritation at the site of product application such as redness, scalaring or other signs of discomfort. If signs of sensitivity occur, bettle your dog with a mild scap and rinse with large amounts of water. If these or other side effects occur, consult your veterinarian or call 1-800-781-4738. If your dog has an unusual reaction to the initial application, consult a veterinarian before repositing application. The most common signs of incidental ingestion are excessive salvation and forming at the mouth. If these symptoms occur, immediately feed your dog and continue to monitor your dog for the next 24 hours.



DO NOT USE ON CATS. Keep cats away from treated dogs for 24 hours. Contact your veterinarian if applied to a cat or ingested by a cat.

EPA Reg No. 2517-176 EPA Est. No. 2517-NE-02

If you have questions or comments about this product please write SENTRY Consumer Response P.O. Box 540399, Ornaha, NE 68154-0399

Manufactured by: Sergeant's Pet Care Products, Inc. Ornalia, NE 68138

In case of Emergency, call: 1-800-781-4738 Non-Emergency call: 1-800-224-PETS (7387)



P006289-1 ANRO







cool (below 77°F/25°C), dry place inaccessible to children and pets. Do not refrigerate, Protect from direct sunlight.

CONTAINER DISPOSAL: If empty: Non-refillable container. Do not reuse or refill this container. Offer for recycling if available. If partly filled: Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

# SENTRY ADVANCE GUARD2 FOR DOGS

Once per month topical flea treatment for dogs This product is for the treatment and prevention of fleas on dogs

- Controls multi-stages of flea lifecycle
- . Kills fleas within 12 hours of application on dogs

READ ENTIRE LABEL BEFORE EACH USE. USE ONLY ON DOGS AND PUPPIES 7 WEEKS OF AGE OR OLDER.
PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS. CAUTION. Harmful if swallowed. Causes moderate eye initation. Avoid contact with eyes or clothing. Wash hands thoroughly with soap and warm water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet.

HAZARDS TO DOMESTIC ANIMALS. For external use on dogs only. Individual sensitivities, white rare, may occur after using ANY pesticide product for dogs. Do not treat your dog with more than one pesticide product at a time. Over dosing your dog can result in serious illness and even death. Consult a veterinarian before using this product on medicated, debilitated, aged, pregnant or nursing dogs.

#### FIRST AID

If swallowed: • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vorniting unless told to do so by the poison control center or doctor. • No not give anything by mouth to an unconscious persor If In eyes: • Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.

If on skin or clothing: • Take off contaminated dothing, Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice HOT LINE NUMBER: Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact 1-800-781-4738 for emergency medical treatment information.

NOTE TO PHYSICIAN: Treat the patient symptomatically.

## DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. TO PREVENT HARM TO YOU AND YOUR DOG, READ ENTIRE LABEL BEFORE EACH USE. FOLLOW ALL DIRECTIONS AND PRECAUTIONARY STATEMENTS CAREFULLY.

- DO NOT ALLOW CHILDREN TO APPLY PRODUCT.
- Do not use on animals other than dogs.
   Do not allow your dog to ingest this product.
   Do not reapply for 4 weeks.
- Use entire contents of tube on each dog. Do not split one tube between dogs. Do not use multiple
- Weigh your dog to be sure you are applying the right dose formulated for the weight of your dog.
- Separate the treated dog from all other dogs and cats for 24 hours after treatment has been applied.
- . Do not have contact or allow children to have contact with the treated area until completely dry.

- How to Open:
  1. Remove product bubes from the peologye.
  2. Separate one bube from the others. Hold the bube with notched end pointing up and away from the lose and body. Use scissors to cut off the narrow end at the not How to Apply:
- 1. Invertitube over dog and use open end to part dog's hair.

2. Squeeze tube firmity to apply all of the solution to the dog's skin as a spot to the dog's back between the shoulder bladies.
6.00 For Dogs Welghing 11-20 Dis. Do not apply to dogs welghing less than 11 lbs. Apply one tube 1.0 mil. (0.034 fl. oz) directly to skin as a spot to the dog's back between the shoulder. blades

#### FRED JEWCY DE APPLICATION

Use this product monthly for control of flea infestations. Studies show that this product kills fleas within 12 hours of application and lasts for up to 4 weeks. This product kills reinfesting fleas within 2 hours and prevents further infestations of fleas. Apply monthly for flea control. This product is waterproof and remains effective, even after battning, swimming, or exposure to sunlight or rain

Side Effects: Monitor your dog after application. Some dogs may experience temporary startle effects when any product is applied. Dogs may experience some temporary initiation of the ster of product application such as redness, scratching or other signs of discomfort. If signs of sensitivity occur, bathe your doo with a mild soon and nose with large amounts of water. If these or other side effects coord, consult your vetermarian or call 1-800-781-4738. If your dog has an unusual reaction to the initial application, consult a vetermarian before repeating application. The most common signs of moderntal impostion are excessive salvation and foaming at the mouth. If these symptoms occur, immediately feed your dog and continue to monitor your dog for the next 2N hours.



DO NOT USE ON CATS. Keep cats away from treated dogs for 24 hours. Contact your veterinarian if applied to a cat or ingested by a cat.

EPA Reg. No. 2517-176 EPA Est. No. 2517-NE-02

If you have questions or comments about this product please write SENTRY Consumer Response P.O. Box 540399, Omaha, NE 68154-0399

Manufactured by: Sergeant's Pet Care Products, Inc.

In case of Emergency, call: 1-800-781-4738 Non-Emergency call: 1-800-224 PETS (7397)



P006290-1 ANRO







# SENTRY ADVANCE GUARD2 FOR DOGS

Once per month topical flea treatment for dogs This product is for the treatment and prevention of fleas on dogs · Controls multi-stages of flea lifecycle

Kills fleas within 12 hours of application on dogs

READ ENTIRE LABEL BEFORE EACH USE.
USE ONLY ON DOGS AND PUPPIES 7 WEEKS OF AGE OR OLDER.

#### PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS, CAUTION. Harmful if swallowed. Causes moderate eye irritation. Avoid contact with eyes or clothing. Wash hands thoroughly with scop and warm water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet.

HAZARDS TO DOMESTIC ANIMALS. For external use on dogs only. Individual sensitivities while rare, may occur after using ANY pesticide product for dogs Do not treat your dog with more than one pesticide product at a time. Over dosing your dog can result in serious likness and even death. Consult a veterinarian before using this product on medicated, debilitated, aged, pregnant or nursing dogs

FIRST AID

If swallowed: • Call a poison control center or doctor immediately for treatment advice, Have person sip a glass of water if able to swaltow.
 Do not induce vorniting unless told to do so by the poison control center or doctor.
 Do not give anything by mouth to an unconscious person. If In eyes: • Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. . Call a poison control center or ductor for treatment advice. If on skin or clothing: • Take off confarminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes Call a poison control center or doctor for treatment advice HOT LINE NUMBER: Have the product container or label with you when caling a poison control center or doctor or going for treatment. You may also contact 1-800-781-4738 for emergency medical treatment information.

NOTE TO PHYSICIAN: Treat the patient symptomatically.

### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. TO PREVENT HARM TO YOU AND YOUR DOG, READ ENTIRE LABEL BEFORE EACH USE. FOLLOW ALL DIRECTIONS AND PRECAUTIONARY STATEMENTS CAREFULLY.

- . DO NOT ALLOW CHILDREN TO APPLY PRODUCT.
- Do not use on animals other than does.
- Do not allow your dog to ingest this product.
- Do not reapply for 4 weeks
- Use entire contents of tube on each dog. Do not split one tube between dogs
- One of the multiple lubles on one at polying the right dose formulated for the weight of your dog
   Weigh your dog to be sure you are applying the right dose formulated for the weight of your dog
   Separate the treated dog from all other dogs and cats for 24 hours after treatment has been applied.
- . Do not have contact or allow children to have contact with the treated area until completely dry.

#### How to Open:

- Remove product tubes from the package,
   Separate one tube from the others. Hold the tube with notched end pointing up and away from the face and body. Use scissors to cut off the narrow end at the notches along the line How to Apply:

How to Apply:

1. Invert labe over dog and use open end to part dog's hair.

2. Squeeze labe firmly to apply all of the solution to the dog's skin as a spot to the dog's back between the shoulder badden and the shoulder badden and the shoulder badden and the shoulder badden. Apply one table 2 Famil, (0.085 fl ox) directly to skin as a spot to the dog's back between the devoted badden. shoulder blades.

#### FREQUENCY OF APPLICATION

Use this product monthly for control of flea intestations. Studies show that this product kills fleas within 12 hours of application and lasts for up to 4 weeks. This product kills reinfesting fleas within 2 hours and prevents further infestations of fleas. Apply monthly for flea control. This product is waterproof and remains effective, even after bathing, swimming, or exposure to sunlight or rain.

Side Effects: Monitor your dog after application. Some dogs may experience temporary startle effects when any product is applied. Dogs may experience some temporary imitation at the site. of product application such as redness, scratching or other signs of discomfort. If signs of sensitivity occur, bathe your dog with a mild soap and rinse with large amounts of water. If these or other side effects occur, consult your veterinarian or call 1-800-781-4738. If your dog has an unusual reaction to the initial application, consult a veterinarian before repeating application. The most common signs of incidental ingestion are excessive salivation and foaming at the mouth if these symptoms occur, immediately feed your dog and continue to monitor your dog for the next 24 hours

DO NOT USE ON CATS. Keep cats away from treated dogs for 24 hours. Contact your veterinarian if applied to a cat or ingested by a cat.


EPA Reg. No. 2517-176 EPA Est. No. 2517-NE-02

If you have questions or comments about this product please write SENTRY Consumer Response P.O. Box 540399, Omaha, NE 68154-0399

Manufactured by: Sergeant's Pet Care Products, Inc.

Omaha, NE 68138

In case of Emergency, call: 1-800-781-4738 Non-Emergency call: 1-800-224-PETS (7387)



P006291-1 ANRO







## STORAGE AND DISPOSAL

PESTICIDE STORAGE: Do not remove tube from the pack until ready to use, Store only in original container in a cool (below 77°F/25°C), dry place inaccessible to children and pets. Do not refrigerate. Protect from direct sunlight.

CONTAINER DISPOSAL: If empty: Non-refillable container. Do not reuse or refill this container. Offer for recycling if available. If partly filled: Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

## SENTRY ADVANCE GUARD2

Once per month topical fee treatment for doos This product is for the treatment and prevention of fleas on dogs
• Controls multi-stages of flea lifecycle

Kills fleas within 12 hours of application on doos

READ ENTIRE LABEL BEFORE EACH USE. USE ONLY ON DOGS AND PUPPIES 7 WEEKS OF AGE OR OLDER. PRECAUTIONARY STATEMENTS

HAZAROS TO HUMANS, CAUTION. Harmful if swallowed, Causes moderate eye initation Avoid contact with eyes or clothing. Wash hands thoroughly with soap and warm water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet.

HAZARDS TO DOMESTIC ANIMALS. For external use on dogs only. Individual sensitivities, while rare, may occur after using ANY pesticide product for dogs. Do not treat your dog with more than one pesticide product at a time. Over dosing your dog can result in serious illness and even death. Consult a veterinarian before using this product on medicated, debilitated. aged, pregnant or nursing dogs.

FIRST AID

If swallowed: . Call a poison control center or doctor immediately for treatment advice . Have person sip a glass of water if able to swallow. • Do not induce vorniting unless told to do so by the poison control center or doctor. • Do not give anything by mouth to an unconscious person If In eyes: • Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice

If on skin or clothing: • Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice. HOT LINE NUMBER: Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact 1-800-781-4738 for emergency medical treatment information.

NOTE TO PHYSICIAN: Treat the patient symptomatically.

#### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. To prevent harm to you and your dog, read entire label before each use. Follow all directions and precautionary statements carefully.

- DO NOT ALLOW CHILDREN TO APPLY PRODUCT.
- Do not use on animals other than dogs.
- . Do not allow your dog to ingest this product.
- · Do not reapply for 4 weeks
- Use entire contents of tube on each dog. Do not split one tube between dogs. Do not use multiple tubes on one doc
- Weigh your dog to be sure you are applying the right dose formulated for the weight of your dog. Separate the treated dog from all other dogs and cats for 24 hours after treatment has been applied.
- Do not have contact or allow children to have contact with the treated area until completely dry.

#### How to Open:

- 1. Remove product tubes from the package.
- Separate one tube from the others. Hold the tube with notched end pointing up and away
  from the face and body. Use scissors to cut off the narrow end at the notches along the line. How to Apply: 1. Invert tube over dog and use open end to part dog's hair.
- 2. Squeeze tube firmly to apply all of the solution to the dog's skin as a spot to the dog's back between the shoulder blades.

Only For Dogs Weighing 55 lbs. or greater. Do not apply to dogs weighing less than 55 lbs. Do not use two tubes on dogs greater than 55 lbs. Apply one tube 4.0 mL (0.135 fl oz) directly to skin as a spot to the dog's back between the shoulder blades.

#### FREQUENCY OF APPLICATION

Use this product monthly for control of fies infestations. Studies show that this product kills fees within 12 hours of application and lasts for up to 4 weeks. This product kills reintesting fees within 2 hours and prevents further infestations of fleas. Apply monthly for flea control. This product is waterproof and remains effective, even after bathing, swimming, or exposure to sunlight or rain.

Side Effects: Monitor your don after application. Some does may experience temporary startle effects when any product is applied. Dogs may experience some temporary initiation at the site of product application such as redness, scratching or other signs of discomfort. If signs of sensitivity occur, bathe your dog with a mild scop and rinse with large amounts of water. If these or other side effects occur, consult your witerinarian or call 1-800-781-4738. If your dog has an unusual reaction to the initial application, consult a veterinarian before repeating application. The most common signs of incidental ingestion are excessive salvation and foarning at the mouth. If these symptoms occur, immediately feed your dog and continue to monitor your dog for the next 24 hours.



DO NOT USE ON CATS. Keep cats away from treated dogs for 24 hours Contact your veterinarian if applied to a cat or ingested by a cat.


EPA Reg No. 2517-176 EPA Est. No. 2517-NE-02

If you have questions or comments about this product please write SENTRY Consumer Response P.O. Box 540399, Ornaha, NE 68154-0399

Manufactured by: Sergeant's Pet Care Products, Inc.

Omaha NF 68139

In case of Emergency, call: 1-800-781-4738 Non-Emergency call: 1-800-224-PETS (7387)



P006292-1 ANRO



