

For use ONLY on cats & kittens Over 1.5 lbs & over 8 weeks of age

CAT MD MAXIMUM DEFENSE PLUS Flea & Tick Topical

3 MONTH SUPPLY

CAT MD MAXIMUM DEFENSE PLUS Flea & Tick Topical

FIPRONIL 9.8% (S)-METHOPRENE 11.8%

For use ONLY on cats & kittens Over 1.5 lbs & over 8 weeks of age

Kills fleas, flea eggs, flea larvae, ticks & chewing lice

Fast acting Long lasting Waterproof

Kills ticks including those that may transmit Lyme disease

KEEP OUT OF REACH OF CHILDREN CAUTION See Back Panel for Additional Precautionary Statements

Table with 2 columns: Ingredient, Percentage. Includes Fipronil (9.80%), (S)-methoprene (11.80%), OTHER INGREDIENT (78.40%), TOTAL (100.00%).

Compare to FRONTLINE PLUS for Cats, contains the same active ingredients.*



UNCOATED AREA

CAT MD MAXIMUM DEFENSE PLUS Flea & Tick Topical

Net Contents: 0.051 fl oz (1.50 mL) • Three 0.017 fl oz (0.50 mL) applicators

STORAGE AND DISPOSAL

PESTICIDE STORAGE: Do not remove tube from the pack until ready to use. Store in a cool (below 77°F/25°C) dry place inaccessible to children and pets. Do not refrigerate. Protect from direct sunlight. CONTAINER DISPOSAL: If empty: Non-refillable container. Do not reuse or refill this container. Offer for recycling if available. If partly filled: Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.



EPA Reg. No. 2517-135-8474 EPA Est. No. 2517-NE-02 In Case of Emergency, call: 1-877-416-7330, Non-Emergency, call: 1-888-814-4738 (4PET)

Distributed by: Merial Animal Health, Omaha, NE 68154 If you have questions or comments about this product, please write: Consumer Response, P.O. Box 540224, Omaha, NE 68154-0224. reply for 30 days.

Use this product monthly for complete control of flea, tick and chewing lice infestations. Studies show that this product kills adult fleas for up to six weeks. Apply monthly if your cat is at high risk for flea infestation. Apply monthly to control ticks and chewing lice. This product remains effective, even after bathing, water immersion, or exposure to sunlight. Allow treated area to dry thoroughly. Do not bathe your cat for 30 days. While holding the cat with one hand, use the other hand to apply the solution. Invert the tube and use the narrow end to part the cat's hair. Apply contents of one tube in paper and put into the trash. Only one applicator per treatment is needed.

How to apply: While holding the cat with one hand, use the other hand to apply the solution. Invert the tube and use the narrow end to part the cat's hair. Apply contents of one tube in paper and put into the trash. Only one applicator per treatment is needed. How to open: Remove product tubes from the package. Separate one tube from the others. Hold the tube with notched end pointing up and away from the face and body. Use scissors to cut off the narrow end at the notches along the line. Side Effects: Monitor your cat after application. Some cats may experience temporary stinging effects when any product is applied. Cats may experience some temporary irritation at the site of product application such as redness, scratching or other signs of discomfort. If signs of sensitivity occur, bathe your cat with a mild soap and rinse with large amounts of water. If these or other side effects occur, consult your veterinarian or call 1-877-416-7330. If your cat has an unusual reaction to the initial application, consult a veterinarian before reapplying. The most common signs of incidental ingestion are excessive salivation and foaming at the mouth. If these symptoms occur, immediately feed your cat and continue to monitor your cat for the next 24 hours.

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. DO NOT ALLOW CHILDREN TO APPLY PRODUCT. TO PREVENT HARM TO YOU AND YOUR CAT READ THE ENTIRE LABEL BEFORE EACH USE. FOLLOW ALL DIRECTIONS AND PRECAUTIONARY STATEMENTS CAREFULLY. Do not use on rabbits or animals other than cats. Do not allow your cat to ingest this product. Do not use on kittens under 8 weeks of age. Do not apply to cats weighing less than 1.5 lbs. Do not double-dose cats. Use entire contents of tube on each cat. Do not split one tube between two cats. Separate the treated cat from all other cats and dogs for 24 hours after treatment has been applied.

DIRECTIONS FOR USE Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-877-416-7330 for emergency medical treatment information.

Table with 2 columns: Action, Instructions. Includes: If swallowed: Call a poison control center or doctor immediately for treatment advice. If on skin: Wash with plenty of soap and water. Get medical attention if irritation persists. If in eyes: Flush eyes with plenty of water. Call a doctor if irritation persists.

FIRST AID Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a healthcare professional. HAZARDS TO HUMANS, CAUTION: Harmful if swallowed. Causes eye irritation. Avoid contact with skin, eyes or clothing. Wash hands thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. HAZARDS TO DOMESTIC ANIMALS: For external use on cats only. Individual sensitivities, while rare, may occur after using any pesticide product on cats. Certain medications can interact with pesticides. Do not treat your cat with more than one pesticide product at a time. Over dosing your cat can result in serious illness and even death. Consult a veterinarian before using on medicated, debilitated, or aged cats. Can be used on breeding, pregnant, or lactating cats. Consult a veterinarian before using on cats with known organ dysfunction.

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS, CAUTION: Harmful if swallowed. Causes eye irritation. Avoid contact with skin, eyes or clothing. Wash hands thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. HAZARDS TO DOMESTIC ANIMALS: For external use on cats only. Individual sensitivities, while rare, may occur after using any pesticide product on cats. Certain medications can interact with pesticides. Do not treat your cat with more than one pesticide product at a time. Over dosing your cat can result in serious illness and even death. Consult a veterinarian before using on medicated, debilitated, or aged cats. Can be used on breeding, pregnant, or lactating cats. Consult a veterinarian before using on cats with known organ dysfunction.

For use ONLY on cats & kittens Over 1.5 lbs & over 8 weeks of age CAT MD MAXIMUM DEFENSE PLUS Flea & Tick Topical

READ ENTIRE LABEL BEFORE EACH USE USE ONLY ON CATS AND KITTENS OVER 8 WEEKS OF AGE CAT MD MAXIMUM DEFENSE PLUS