

APPROVED

By Lisa Adamson at 11:49 am, Nov 24, 2014

ZoGuard plus FOR CATS

ZoGuard Plus for Cats contains Fipronil and S-Methoprene (insect growth regulator) to effectively kill fleas (all stages) and ticks (including those that may transmit Lyme disease). The active ingredients in ZoGuard Plus for Cats feature a sustained release formula developed for fast-acting and long-lasting control of fleas, ticks, and chewing lice for 1 month on cats only.

DUAL ACTION!

Fipronil with (S)-methoprene IGR Effectively Breaks the Flea Life Cycle



FLEA EGGS FLEA LARVAE FLEA PUPAE ADULT FLEAS

- Use ZoGuard Plus for Cats monthly to break the flea life cycle.
- Kills fleas, ticks, chewing lice, flea larvae & prevents the development of flea pupae
- Kills Ticks including those that may transmit Lyme disease
- Kills brown dog ticks, American dog ticks, lone star ticks, and deer ticks for up to 1 month
- Rapid elimination of chewing lice infestation
- Fast-acting
- Long-lasting
- Waterproof
- Can be used on breeding, pregnant and lactating cats

STORAGE AND DISPOSAL: Do not remove tube from the pack until ready to use. Store in a cool, dry place (below 77°F / 25°C) dry place inaccessible to children and pets. Do not refrigerate. Protect from direct sunlight. Do not use or refill this container. Offer for recycling if available. If readily filled, call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

ONLY for cats & kittens over 8 weeks of age and over 1.5 lbs

In Case of Emergency, call: 1-844-685-9173

Made in USA

READ ENTIRE LABEL BEFORE EACH USE. USE ONLY ON CATS AND KITTENS OVER 8 WEEKS OF AGE.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS. CAUTION. Harmful if swallowed. Causes eye irritation. Avoid contact with skin, eyes or clothing. Wash hands thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet.

HAZARDS TO DOMESTIC ANIMALS. For external use on cats only. Individual sensitivities, while rare, may occur after using any pesticide product on cats. Certain medications can interact with pesticides. Do not treat your cat with more than one pesticide product at a time. Over dosing your cat can result in serious illness and even death. Consult a veterinarian before using on medicated, debilitated, or aged cats. Can be used on breeding, pregnant, or lactating cats. Consult a veterinarian before using on cats with known organ dysfunction.

FIRST AID

If swallowed:	Call a poison control center or a doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.
If in eyes:	Flush eyes with plenty of water. Call a doctor if irritation persists.
If on skin:	Wash with plenty of soap and water. Get medical attention if irritation persists.

HOT LINE NUMBER

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-844-685-9173 for emergency medical treatment information.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. **DO NOT ALLOW CHILDREN TO APPLY PRODUCT.** TO PREVENT HARM TO YOU AND YOUR CAT, READ ENTIRE LABEL BEFORE EACH USE. FOLLOW ALL DIRECTIONS AND PRECAUTIONARY STATEMENTS CAREFULLY. Do not use on rabbits or animals other than cats. Do not allow your cat to ingest this product. Do not use on kittens under 8 weeks of age. Do not apply to cats weighing less than 1.5 lbs. Do not double-dose cats. Use entire contents of tube vial on each cat. Separate the treated cat from all other cats and dogs for 24 hours after treatment has been applied.

Side Effects: Monitor your cat after application. Cats may experience some temporary irritation at the site of product application such as redness, scratching or other signs of discomfort. If signs of sensitivity occur, bathe your cat with a mild soap and rinse with large amounts of water. If these or other side effects occur consult your veterinarian or call 1-844-685-9173. If your cat has an unusual reaction to the initial application, consult a veterinarian before repeating application. The most common signs of incidental ingestion are excessive salivation and foaming at the mouth. If these symptoms occur, immediately feed your cat and continue to monitor your cat for the next 24 hours.

How to open: Remove product tubes from the package. Separate one tube from the others. Hold the tube with notched end pointing up and away from the face and body. Use scissors to cut off the narrow end at the notches along the lines.

How to apply: While holding the cat with one hand, use the other hand to apply the solution. Invert the tube and use the narrow end to part the cat's hair. Apply contents of one tube of ZoGuard Plus for Cats as a spot between the shoulder blades. Hold the cat for a few seconds to give the solution time to be absorbed into the cat's coat. When finished, wrap the tube in paper and put into the trash. Only one applicator per treatment is needed.

FREQUENCY OF APPLICATION

Use ZoGuard™ Plus for Cats monthly for complete control of flea, tick, and chewing lice infestations. Studies show that ZoGuard™ Plus for Cats kills adult fleas for up to six weeks. Apply monthly if your cat is at high risk for flea reinfestation. Apply monthly to control ticks and chewing lice. ZoGuard™ Plus for Cats remains effective even after bathing, water immersion, or exposure to sunlight. Allow treated area to dry thoroughly. Do not reapply for 30 days.

If you have questions or comments about this product, please write:

Promika, LLC
1204 Village Market Pl #273, Morrisville, NC 27560

Distributed by:
Promika, LLC
1204 Village Market Place #273
Morrisville, NC 27560

EPA REG. NO. 53883-352-91300
EPA EST. NO. 2517-NE-001

ZoGuard plus FOR CATS

ONLY for cats & kittens over 8 weeks of age and over 1.5 lbs
P005924



3 MONTH SUPPLY

3 MONTH SUPPLY

ONLY for cats & kittens over 8 weeks of age and over 1.5 lbs

ZoGuard plus FOR CATS



Contains Fipronil and S-Methoprene, the active ingredients used in FRONTLINE® Plus for Cats

- FLEAS
- TICKS
- FLEA EGGS
- FLEA LARVAE
- CHEWING LICE

KEEP OUT OF REACH OF CHILDREN

CAUTION See Back Panel For Additional Precautionary Statements.

NET CONTENTS: 0.051 fl oz
Three 0.017 fl oz (0.50 mL) Applicators
Frontline® is a registered trademark of Merial. ZoGuard™ Plus for Cats is not manufactured, or distributed by Merial, seller of Frontline® Plus for Cats.

ACTIVE INGREDIENTS:
Fipronil: 9.800%
(S)-methoprene: 11.800%
OTHER INGREDIENTS: 78.400%
TOTAL: 100.00%

LIFT HERE FOR MORE INFORMATION

DIRECTIONS FOR USE


How to Open

1. Remove product tubes from the package.
2. Separate one tube from the others. Hold the tube with notched end pointing up and away from the face and body. Use scissors to cut off the narrow end at the notches along the line.



How to Apply

1. Invert tube over cat and use open end to part cat's hair.
2. Apply contents of one tube of ZoGuard Plus for Cats as a spot between the shoulder blades. Squeeze tube firmly to apply all of the solution to the cat's skin.



6 STEP CHECKLIST

1. Read the label completely and follow directions
2. Weigh your cat
3. Do NOT use dog products on cats
4. Do NOT treat with more than one pesticide product at a time
5. Separate animals after treatment to avoid chance of ingestion
6. Do NOT split tubes between pets

3 MONTH TRACKER

Treatment 1	MM	DD	YY
Treatment 2	MM	DD	YY
Treatment 3	MM	DD	YY