

For use ONLY on dogs & puppies 4-22 lbs & over 8 weeks of age

DOG MD
MAXIMUM DEFENSE™ **PLUS**
Flea & Tick Topical

3 MONTH SUPPLY

DOG MD

MAXIMUM DEFENSE™ PLUS

Flea & Tick Topical

3 MONTH SUPPLY

Compare to FRONTLINE® PLUS for Dogs, contains the same active ingredients.*

FIPRONIL 9.8%
(S)-METHOPRENE 8.8%

For use ONLY on dogs & puppies 4-22 lbs & over 8 weeks of age

Kills fleas, flea eggs, flea larvae, ticks & chewing lice

Fast acting
Long lasting
Waterproof

Kills ticks including those that may transmit Lyme disease



KEEP OUT OF REACH OF CHILDREN CAUTION

See Back Panel for Additional Precautionary Statements



DO NOT USE ON CATS

ACTIVE INGREDIENTS:


Fipronil	9.80%
(S)-methoprene	8.80%
OTHER INGREDIENT:	81.40%
TOTAL	100.00%

*This product is not manufactured or distributed by Merial, the makers of FRONTLINE® Plus for Dogs. FRONTLINE® is a registered trademark of Merial.

Net Contents: 0.069 fl oz (2.01 mL) • Three 0.023 fl oz (0.67 mL) applicators

STORAGE AND DISPOSAL

PESTICIDE STORAGE: Do not remove tube from the pack until ready to use. Store in a cool (below 77°F/25°C) dry place inaccessible to children and pets. Do not refrigerate. Protect from direct sunlight.
CONTAINER DISPOSAL: If empty: Non-refillable container. Do not reuse or refill this container. Offer for recycling if available. If partly filled: Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.



DO NOT USE ON CATS. Keep cats away from treated dogs for 24 hours. If applied to a cat or ingested by a cat, contact your veterinarian.

How to open: Remove product tubes from the package. Separate one tube from the others. Hold the tube with notched end pointing up and away from the face and body. Use scissors to cut off the narrow end at the notches along the line.
How to apply: Invert tube over dog and use open end to part dog's hair. Squeeze tube firmly to apply all of the solution to the dog's skin, as directed below. Do not apply more often than once every 4 weeks. **Only for Dogs Weighing 4-22 lbs. Do not apply to dogs weighing less than 4 lbs.** Apply one tube (0.67 mL) (0.023 fl oz) as a spot to the dog's back between the shoulder blades.
FREQUENCY OF APPLICATION
Use this product monthly for complete control of flea, tick, and chewing lice infestations. Studies show that this product kills adult fleas for up to three months. Apply monthly for flea reinfestation, or exposure to sunlight, allow treated area to dry thoroughly. Do not reapply for 30 days.
If you have questions or comments about this product, please write: Consumer Response, P.O. Box 540224, Omaha, NE 68154-0224.
Distributed by: Meridian Animal Health, Omaha, NE 68154
In Case of Emergency, call: 1-877-416-7330, Non-Emergency, call: 1-888-814-4738 (4PET)

Side Effects: Monitor your dog after application. Some dogs may experience temporary startle effects when any product is applied. Dogs may experience some temporary irritation at the site of product application such as redness, scratching or other signs of discomfort. If signs of sensitivity occur, bathe your dog with a mild soap and rinse with large amounts of water. If these or other side effects occur, consult your veterinarian or call 1-877-416-7330. If your dog has an unusual reaction to the initial application, consult a veterinarian before reapplying. The most common signs of incidental ingestion are excessive salivation and foaming at the mouth. If these symptoms occur, immediately feed your dog and continue to monitor your dog for the next 24 hours.

First Aid:
If swallowed: Call a poison control center or a doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.
If in eyes: Flush eyes with plenty of water. Call a doctor if irritation persists.
If on skin: Wash with plenty of soap and water. Get medical attention if irritation persists.

Precautionary Statements:
Hazardous to humans, CAUTION: Harmful if swallowed or using the toilet.
Avoid contact with skin, eyes or clothing. Wash hands thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet.
Hazardous to domestic animals. For external use on dogs only. Individual sensitivities, while rare, may occur after using any pesticide product on dogs. Certain medications can interact with pesticides. Do not treat your dog with more than one pesticide product at a time. Over dosing your dog can result in serious illness and even death. Consult a veterinarian before using on medicated, dehydrated, aged, pregnant or nursing dogs. Consult a veterinarian before using on dogs with known organ dysfunction. If your dog is exhibiting signs of and/or is being treated for skin dermatitis, talk to your veterinarian before applying any topical flea and tick control product.

READ ENTIRE LABEL BEFORE EACH USE. USE ONLY ON DOGS AND PUPPIES OVER 8 WEEKS OF AGE.

DOG MD
MAXIMUM DEFENSE™ **PLUS**
Flea & Tick Topical

For use ONLY on dogs & puppies & over 8 weeks of age 4-22 lbs

DOG MD
MAXIMUM DEFENSE™ **PLUS**
Flea & Tick Topical

UNCOATED AREA

DOG MD
MAXIMUM DEFENSE™ **PLUS**
Flea & Tick Topical

For use ONLY on dogs **23-44 lbs**

DOGMD
MAXIMUM DEFENSE™ **PLUS**
Flea & Tick Topical

3 MONTH SUPPLY

DOGMD

MAXIMUM DEFENSE™ **PLUS**

Flea & Tick Topical

3 MONTH SUPPLY

Compare to **FRONTLINE® PLUS** for Dogs, contains the same active ingredients.*

FIPRONIL 9.8%
(S)-METHOPRENE 8.8%

For use ONLY on dogs **23-44 lbs**

Kills fleas, flea eggs, flea larvae, ticks & chewing lice

Fast acting
Long lasting
Waterproof

Kills ticks including those that may transmit Lyme disease

KEEP OUT OF REACH OF CHILDREN
CAUTION See Back Panel for Additional Precautionary Statements



DO NOT USE ON CATS

ACTIVE INGREDIENTS:
Fipronil.....9.80%
(S)-methoprene.....8.80%
OTHER INGREDIENTS:.....81.40%
TOTAL.....100.00%

*This product is not manufactured or distributed by Merial, the makers of FRONTLINE® Plus for Dogs. FRONTLINE® is a registered trademark of Merial.

Net Contents: 0.135 fl oz (4.02 mL) • Three 0.045 fl oz (1.34 mL) applicators

STORAGE AND DISPOSAL

PESTICIDE STORAGE: Do not remove tube from the pack until ready to use. Store in a cool (below 77°F/25°C), dry place inaccessible to children and pets. Do not refrigerate. Protect from direct sunlight.
CONTAINER DISPOSAL: If empty: Non-refillable container. Do not reuse or refill this container. Offer for recycling if available. If partly filled: Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.



EPA Reg. No. 2517-134-9474Z EPA Est. No. 2517-1A-02

In Case of Emergency, call: 1-877-416-7330, Non-Emergency, call: 1-888-814-4738 (4PET)
Distributed by: Meridian Animal Health, Omaha, NE 68154
If you have questions or comments about this product, please write: Consumer Response, P.O. Box 540224, Omaha, NE 68154-0224.
not reapply for 30 days.

This product remains effective even after bathing, water immersion, or exposure to sunlight. Allow treated area to dry thoroughly. Do not apply to wet fur.
Use this product monthly for complete control of flea, tick, and chewing lice infestations. Studies show that this product kills adult fleas on up to three months. Apply monthly for flea reinfestation. Apply monthly to control ticks and chewing lice.

FREQUENCY OF APPLICATION

How to open: Remove product tubes from the package. Separate one tube from the others. Hold the tube with notched end pointing up and away from the face and body. Use scissors to cut off the narrow end at the notches along the line.

How to apply: Invert tube over dog and use open end to part dog's hair. Squeeze tube firmly to apply all of the solution to the dog's skin, as directed below. Do not apply to the face and neck area. Apply one tube (1.34 mL/0.045 fl oz) as a spot to the dog's back between the shoulder blades.

DO NOT USE ON CATS. Keep cats away from treated dogs for 24 hours. If applied to a cat or ingested by a cat, contact your veterinarian.
Site of product application: Monitor your dog after application. Some dogs may experience temporary startle effects when any product is applied. Dogs may experience some temporary irritation at the site of product application such as redness, soreness, or other signs of discomfort. If signs of sensitivity occur, bathe your dog with a mild soap and rinse with large amounts of water; if these or other side effects occur, consult your veterinarian or call 1-877-416-7330. If your dog has an unusual reaction to the initial application, consult a veterinarian before reapplying application. The most common signs of incidental ingestion are excessive salivation and foaming at the mouth. If these symptoms occur, immediately feed your dog and continue to monitor your dog for the next 24 hours.

HAZARDS TO DOMESTIC ANIMALS: For external use on dogs only. Individual sensitivities, while rare, may occur after using any pesticide product on dogs. Certain medications can interact with pesticides. Do not treat your dog with more than one pesticide product at a time. Over dosing your dog can result in serious illness and even death. Consult a veterinarian before eating, drinking, chewing gum, using tobacco or using the toilet.
HAZARDS TO HUMANS. CAUTION. Harmful if swallowed. Causes eye irritation. Avoid contact with skin, eyes or clothing. Wash hands thoroughly with soap and water after handling and is being treated for skin dermatitis, talk to your veterinarian before applying any topical flea and tick control product.

READ ENTIRE LABEL BEFORE EACH USE. FOLLOW ALL DIRECTIONS AND PRECAUTIONARY STATEMENTS CAREFULLY. Do not use on cats, rabbits or animals other than dogs. Do not allow your dog to ingest this product. Do not use on puppies under 8 weeks of age or less than 23 lbs. Use entire contents of tube on each dog. Do not split one tube between dogs. Do not use multiple tubes on one dog. Weigh your dog to be sure you are applying the right dose formulated for the weight of your dog. Separate the treated dog from all other dogs and cats for 24 hours after treatment has been applied.

IF SWALLOWED: Call a poison control center or a doctor immediately for treatment advice. Have person sip glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.
IF IN EYES: Flush eyes with plenty of water. Call a doctor if irritation persists.
IF ON SKIN: Wash with plenty of soap and water. Get medical attention if irritation persists.

HOT LINE NUMBER
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-877-416-7330 for emergency medical treatment information.

FIRST AID
Call a poison control center or a doctor immediately for treatment advice. Have person sip glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.

PRECAUTIONARY STATEMENTS
USE ONLY ON DOGS AND PUPPIES OVER 8 WEEKS OF AGE.
READ ENTIRE LABEL BEFORE EACH USE.

DOGMD
MAXIMUM DEFENSE™ **PLUS**
Flea & Tick Topical

For use ONLY on dogs **23-44 lbs**

DOGMD
MAXIMUM DEFENSE™ **PLUS**
Flea & Tick Topical

UNCOATED AREA

DOGMD
MAXIMUM DEFENSE™ **PLUS**
Flea & Tick Topical

For use ONLY on dogs 45-88 lbs

DOGMD
MAXIMUM DEFENSE™ **PLUS**
Flea & Tick Topical

3 MONTH SUPPLY

DOGMD

MAXIMUM DEFENSE™ **PLUS**
Flea & Tick Topical

3 MONTH SUPPLY

Compare to FRONTLINE® PLUS for Dogs, contains the same active ingredients.*

FIPRONIL 9.8%
(S)-METHOPRENE 8.8%

For use ONLY on dogs 45-88 lbs

Kills fleas, flea eggs, flea larvae, ticks & chewing lice

Fast acting
Long lasting
Waterproof

Kills ticks including those that may transmit Lyme disease



KEEP OUT OF REACH OF CHILDREN
CAUTION See Back Panel for Additional Precautionary Statements



DO NOT USE ON CATS

ACTIVE INGREDIENTS:
Fipronil 9.80%
(S)-methoprene 8.80%
OTHER INGREDIENT: 81.40%
TOTAL 100.00%

*This product is not manufactured or distributed by Merial, the makers of FRONTLINE® Plus for Dogs. FRONTLINE® is a registered trademark of Merial.

Net Contents: 0.273 fl oz (8.04 mL) • Three 0.091 fl oz (2.68 mL) applicators

STORAGE AND DISPOSAL

PESTICIDE STORAGE: Do not remove tube from the pack until ready to use. Store in a cool (below 77°F/25°C), dry place inaccessible to children and pets. Do not refrigerate. Protect from direct sunlight.
CONTAINER DISPOSAL: If empty: Non-refillable container. Do not reuse or refill this container. Offer for recycling if available. If partly filled: Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.



DO NOT USE ON CATS. Keep cats away from treated dogs for 24 hours. If applied to a cat or ingested by a cat, contact your veterinarian.



It is a violation of Federal law to use this product in a manner inconsistent with its labeling. **DO NOT ALLOW CHILDREN TO APPLY PRODUCT.** TO PREVENT HARM TO YOU AND YOUR DOG, READ THE LABEL BEFORE EACH USE. FOLLOW ALL DIRECTIONS AND PRECAUTIONARY STATEMENTS CAREFULLY. Do not use on cats, rabbits or animals other than dogs. Do not allow multiple tubes on one dog. Weight your dog to be sure you are applying the right dose formulated for the weight of your dog. Separate the treated dog from all other dogs and cats for 24 hours after treatment has been applied.

SIDE EFFECTS: Monitor your dog after application. Some dogs may experience temporary startle effects when any product is applied. Dogs may experience some temporary irritation at the site of product application such as redness, scratching or other signs of discomfort. If signs of sensitivity occur, bathe your dog with a mild soap and rinse with large amounts of water. If these or other side effects occur, consult your veterinarian or call 1-877-416-7330. If your dog has an unusual reaction to the initial application, consult a veterinarian before repeating application. The most common signs of incidental ingestion are excessive salivation and foaming at the mouth. If these symptoms occur, immediately feed your dog and continue to monitor your dog for the next 24 hours.

DIRECTIONS FOR USE	
If swallowed:	Call a poison control center or a doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or a doctor. Do not give anything by mouth to an unconscious person.
If in eyes:	Flush eyes with plenty of water. Call a doctor if irritation persists.
If on skin:	Wash with plenty of soap and water. Get medical attention if irritation persists.
HOT LINE NUMBER	
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-877-416-7330 for emergency medical treatment information.	

FIRST AID

Call a poison control center or a doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or a doctor. Do not give anything by mouth to an unconscious person.

If in eyes: Flush eyes with plenty of water. Call a doctor if irritation persists.

If on skin: Wash with plenty of soap and water. Get medical attention if irritation persists.

HAZARDS TO HUMANS, CAUTION. Harmful if swallowed. Causes eye irritation. Avoid contact with skin, eyes or clothing. Wash hands thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet.

HAZARDS TO DOMESTIC ANIMALS. For external use on dogs only. Individual sensitivities, while rare, may occur after using any pesticide product on dogs. Certain medications can interact with pesticides. Do not treat your dog with more than one pesticide product at a time. Over dosing your dog can result in serious illness and even death. Consult a veterinarian before using on medicated, debilitated, aged, pregnant or nursing dogs. Consult a veterinarian before applying any topical flea and tick control product.

As being treated for skin dermatitis, talk to your veterinarian before applying any topical flea and tick control product.

READ ENTIRE LABEL BEFORE EACH USE. USE ONLY ON DOGS AND PUPPIES OVER 8 WEEKS OF AGE. PRECAUTIONARY STATEMENTS

For use ONLY on dogs 45-88 lbs

DOGMD
MAXIMUM DEFENSE™ **PLUS**
Flea & Tick Topical

UNCOATED AREA

DOGMD
MAXIMUM DEFENSE™ **PLUS**
Flea & Tick Topical

For use ONLY on dogs 89-132 lbs

DOGMD
MAXIMUM DEFENSE™ **PLUS**
Flea & Tick Topical

3 MONTH SUPPLY

DOGMD

MAXIMUM DEFENSE™ **PLUS**
Flea & Tick Topical

3 MONTH SUPPLY

Compare to FRONTLINE® PLUS for Dogs, contains the same active ingredients.*

FIPRONIL 9.8%
(S)-METHOPRENE 8.8%

For use ONLY on dogs 89-132 lbs

Kills fleas, flea eggs, flea larvae, ticks & chewing lice

Fast acting
Long lasting
Waterproof

Kills ticks including those that may transmit Lyme disease

KEEP OUT OF REACH OF CHILDREN
CAUTION See Back Panel for Additional Precautionary Statements



DO NOT USE ON CATS

ACTIVE INGREDIENTS:
Fipronil.....9.80%
(S)-methoprene.....8.80%
OTHER INGREDIENT:.....81.40%
TOTAL.....100.00%

*This product is not manufactured or distributed by Merial, the makers of FRONTLINE® Plus for Dogs. FRONTLINE® is a registered trademark of Merial.

Net Contents: 0.408 fl oz (12.06 mL) • Three 0.136 fl oz (4.02 mL) applicators

STORAGE AND DISPOSAL

PESTICIDE STORAGE: Do not remove tube from the pack until ready to use. Store in a cool (below 77°F/25°C), dry place inaccessible to children and pets. Do not refrigerate. Protect from direct sunlight.
CONTAINER DISPOSAL: If empty: Non-refillable container. Do not reuse or refill this container. Offer for recycling if available. If partly filled: Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.



FREQUENCY OF APPLICATION
Use this product monthly for complete control of flea, tick, and chewing lice infestations. Studies show that this product kills adult fleas for up to three months. Apply monthly if your dog is at high risk for flea reinfestation. Apply monthly to control ticks and chewing lice. This product remains effective even after bathing, water immersion, or exposure to sunlight. Allow treated area to dry thoroughly. Do not reapply for 30 days.
If you have questions or comments about this product, please write: Consumer Response, P.O. Box 540224, Omaha, NE 68154-0224.
Distributed by: Meridian Animal Health, Omaha, NE 68154
In Case of Emergency, call: 1-877-416-7330, Non-Emergency, call: 1-888-814-4738 (4PET)



DO NOT USE ON CATS. Keep cats away from treated dogs for 24 hours. If applied to a cat or ingested by a cat, contact your veterinarian.
How to open: Remove product tubes from the package. Separate one tube from the others. Hold the tube with notched end pointing up and away from the face and body. Use scissors to cut off the narrow end at the notches along the line.
How to apply: Invert tube over dog and use open end to part dog's hair. Squeeze tube firmly to apply all of the solution to the dog's skin, as directed below. Do not apply more often than once every 4 weeks. **Only for Dogs Weighing 89-132 lbs. Do not use two tubes on dogs greater than 132 lbs.** Apply one tube (4.02 mL) (0.136 fl oz) as a spot to the dog's back between the shoulder blades.

Side Effects: Monitor your dog after application. Some dogs may experience temporary startle effects when any product is applied. Dogs may experience some temporary irritation at the site of product application such as redness, scratching or other signs of discomfort. If signs of sensitivity occur, bathe your dog with a mild soap and rinse with large amounts of water. If these or other side effects occur, consult your veterinarian or call 1-877-416-7330. If your dog has an unusual reaction to the initial application, consult a veterinarian before reapplying. application. The most common sign of incidental ingestion are excessive salivation and foaming at the mouth. If these symptoms occur, immediately feed your dog and confine to monitor your dog for the next 24 hours.

READ ENTIRE LABEL BEFORE EACH USE. FOLLOW ALL DIRECTIONS AND PRECAUTIONARY STATEMENTS CAREFULLY. Do not use on cats, rabbits or animals other than dogs. Do not allow multiple tubes on one dog. Weigh your dog to be sure you are applying the right dose formulated for the weight of your dog. Separate the treated dog from all other dogs and cats for 24 hours after treatment has been applied.

READ ENTIRE LABEL BEFORE EACH USE. FOLLOW ALL DIRECTIONS AND PRECAUTIONARY STATEMENTS CAREFULLY. Do not use on cats, rabbits or animals other than dogs. Do not allow multiple tubes on one dog. Weigh your dog to be sure you are applying the right dose formulated for the weight of your dog. Separate the treated dog from all other dogs and cats for 24 hours after treatment has been applied.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-877-416-7330 for emergency medical treatment information.

HOT LINE NUMBER

If on skin: Wash with plenty of soap and water. Get medical attention if irritation persists.
If in eyes: Flush eyes with plenty of water. Call a doctor if irritation persists.
If swallowed: Call a poison control center or a doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.

FIRST AID

HAZARDS TO DOMESTIC ANIMALS: For external use on dogs only. Individual sensitivities, while rare, may occur after using any pesticide product on dogs. Certain medications can interact with pesticides. Do not treat your dog with more than one pesticide product at a time. Over dosing your dog can result in serious illness and even death. Consult a veterinarian before using a pesticide product on dogs with known organ dysfunction. If your dog is exhibiting signs of and/or before using a pesticide product on dogs with known organ dysfunction. If your dog is exhibiting signs of and/or

HAZARDS TO HUMANS, CAUTION. Harmful if swallowed. Causes eye irritation. Avoid contact with skin, eyes or clothing. Wash hands thoroughly with soap and water after handling and

PRECAUTIONARY STATEMENTS
USE ONLY ON DOGS AND PUPPIES OVER 8 WEEKS OF AGE.
READ ENTIRE LABEL BEFORE EACH USE.

For use ONLY on dogs 89-132 lbs

DOGMD
MAXIMUM DEFENSE™ **PLUS**
Flea & Tick Topical

UNCOATED AREA

DOGMD
MAXIMUM DEFENSE™ **PLUS**
Flea & Tick Topical

